

Boğaziçi MATH GRAD SEMINAR

Trace Formula for Finite Groups

Meleknaz Uzuner
Boğaziçi University

Abstract:

Let G be a finite group and consider a subgroup Γ of G along with a representation $\rho : \Gamma \rightarrow \text{GL}(V_\rho)$ of Γ on a finite dimensional \mathbb{C} -linear vector space V_ρ . We get the trace formula by computing the trace of the operator $\text{Ind}_\Gamma^G(\rho)(f) : \text{Ind}_\Gamma^G(V_\rho) \rightarrow \text{Ind}_\Gamma^G(V_\rho)$ for a test function $f : G \rightarrow \mathbb{C}$. Our aim is to present this computation in two different ways, that is, by expressing the spectral side $J(\rho, f)$ and the geometric side $I(\rho, f)$ of the formula. The expression $I(\rho, f)$ is a sum over non-isomorphic irreducible representations of G having the multiplicities of the irreducible representations emerging in the decomposition of the induced representation $\text{Ind}_\Gamma^G(\rho)$ within and the geometric side $I(\rho, f)$ of the trace formula is obtained by calculating the conjugacy classes of the groups Γ and G . We call the generalized formula $J(\rho, f) = \text{Tr}(\text{Ind}_\Gamma^G(\rho)(f)) = I(\rho, f)$ the trace formula for the finite group G with respect to the subgroup Γ and $\rho : \Gamma \rightarrow \text{GL}(V_\rho)$.

Date : Tuesday, May 13, 2025

Time: 13:30

Place: TB130, South Campus, Boğaziçi University